

# PEDIATRICS®

OFFICIAL JOURNAL OF THE AMERICAN ACADEMY OF PEDIATRICS

## **Cultural Bias and Circumcision: The AAP Task Force on Circumcision Responds**

### **TASK FORCE ON CIRCUMCISION**

*Pediatrics*; originally published online March 18, 2013;  
DOI: 10.1542/peds.2013-0081

The online version of this article, along with updated information and services, is  
located on the World Wide Web at:  
<http://pediatrics.aappublications.org/content/early/2013/03/12/peds.2013-0081.citation>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2013 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



# Cultural Bias and Circumcision: The AAP Task Force on Circumcision Responds

**AUTHOR:** TASK FORCE ON CIRCUMCISION

**KEY WORDS**

circumcision, HIV, urinary tract infection, sexually transmitted infections

www.pediatrics.org/cgi/doi/10.1542/peds.2013-0081

doi:10.1542/peds.2013-0081

Accepted for publication Jan 17, 2013

Address correspondence to Douglas S. Diekema, MD, MPH, Seattle Children's Research Institute, C9S-6, 1900 Ninth Ave, Seattle, WA 98101. E-mail: diek@u.washington.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2013 by the American Academy of Pediatrics

**FINANCIAL DISCLOSURE:** *Dr Carlo is a Director of Mednax; the other authors have indicated they have no financial relationships relevant to this article to disclose.*

**FUNDING:** No external funding.

**COMPANION PAPER:** A companion to this article can be found on page 796, and online at [www.pediatrics.org/cgi/doi/10.1542/peds.2012-2896](http://www.pediatrics.org/cgi/doi/10.1542/peds.2012-2896).

FREE

The members of the American Academy of Pediatrics (AAP) Task Force on Circumcision appreciate the opportunity to respond to the concerns raised by Frisch et al in their commentary, "Cultural Bias in AAP's 2012 Technical Report and Policy Statement on Male Circumcision."

The central claim of these authors is that the conclusions of the task force report are culturally biased, leading the task force to a flawed understanding of what constitutes trustworthy evidence and to conclusions that are far from those reached by physicians in most other Western countries. The "obvious" cultural bias to which they refer apparently has its genesis in "the normality of non-therapeutic male circumcision in the US." All of the commentary authors hail from Europe, where the vast majority of men are uncircumcised and the cultural norm clearly favors the uncircumcised penis. In contrast, approximately half of US males are circumcised, and half are not.<sup>1</sup> Although that heterogeneity may lead to a more tolerant view toward circumcision in the United States than in Europe, the cultural "bias" in the United States is much more likely to be a neutral one than that found in Europe, where there is a clear bias against circumcision. Yet, the commentary's authors have, at no point, recognized that their own cultural bias may exist in equal, if not greater, measure than any cultural bias that might exist among the members of the AAP Task Force on Circumcision. If cultural bias influences the review of available evidence, then a culture that is comfortable with both the circumcised penis and the uncircumcised penis would seem predisposed to a more dispassionate analysis of the scientific literature than a culture with a bias that is either strongly opposed to circumcision or strongly in favor of it.

The task force's process was systematic, objective, comprehensive, and transparently documented in its technical report.<sup>1</sup> Members of the AAP Task Force on Circumcision were recruited on the basis of area of expertise. There was no consideration or knowledge of the individuals' beliefs concerning circumcision at the time of their appointment. Unlike other published policy statements and reports on circumcision, the task force did not selectively choose which articles to review, but reviewed all of the available literature identified in a comprehensive search and evaluated those manuscripts by using previously established, nationally recognized guidelines to determine the quality of the data being reviewed.<sup>2</sup> Some articles were reviewed but not cited in the technical report, either because they were not data-based studies, the quality of the study was seriously flawed, or the findings of the study did not meaningfully address the specific area of task force inquiry.

Frisch et al present opinions that reflect a review of the literature that is not comprehensive, systematic, or unbiased. For example, the authors

dismiss the data related to urinary tract infection on the grounds that no randomized controlled trial has been performed, despite the fact that there is good evidence from other studies that suggest a preventive benefit of circumcision. At the same time, they readily dismiss 3 randomized controlled trials and 11 other studies that provide good to fair evidence of a reduction in HIV acquisition associated with circumcision. They claim that these data are “contradicted by other studies, which show no relationship between HIV infection rates and circumcision status,” yet support that claim with only a single reference to a review article authored by the vice president of an organization opposed to circumcision. We would refer the reader to the task force’s technical report for a comprehensive review of the literature related to the potential benefits of circumcision.<sup>1</sup> Notably, the World Health Organization has concluded that the data strongly support a benefit of **male circumcision** with regard to prevention of HIV infection and has issued guidelines for its use, both for adults/adolescents and for neonates.<sup>3</sup>

## COMPLICATIONS

Frisch et al charge that members of the AAP Task Force on Circumcision “consider the foreskin to be a part of the male body that has no meaningful function in sexuality.” They additionally claim, “Recent studies ... suggest that circumcision desensitizes the penis and may lead to sexual problems in circumcised men and their partners.” In fact, many of these studies were reviewed by members of the task force but were not cited in the technical report, either because the findings were equivocal, they did not support a benefit or detriment with regard to sexual function and pleasure, or because the relevance to

individuals undergoing circumcision during infancy was questionable. For example, the authors cite 5 studies to support the claim that “the foreskin is a richly innervated structure that protects the glans and plays an important role in the mechanical function of the penis during sexual acts.” Of these 5 studies, 4 were histologic studies that were not designed to correlate anatomic findings with physiologic or functional roles.<sup>4–7</sup> **Members of the task force appreciate that the foreskin has nerve fibers: the task force clearly recommends adequate pain control for infants undergoing circumcision.** However, the task force did not move beyond what these studies actually reveal (the foreskin has nerve bundles and pain fibers, the foreskin contains Meissner corpuscles, the inner surface of the foreskin resembles a mucous membrane) **to speculate about the effect that circumcision might have on sexual function or pleasure.** The fifth study cited was designed to determine whether the absence of the penile-cavernosus reflex is a reliable indicator of a pathologic sacral lesion and did not evaluate implications for sexual function or pleasure.<sup>8</sup> In sum, of the 5 studies, not one sought to evaluate whether the foreskin protects the glans or whether it “plays an important role in the mechanical function of the penis during sexual acts.”

The authors cite 2 articles as evidence that “circumcision desensitizes the penis.” One of these is not a study and does not present data.<sup>9</sup> The other revealed that perception sensitivity to vibration decreases after circumcision.<sup>10</sup> Neither the clinical implications nor impact on sexual experience were evaluated.

Finally, Frisch et al cite 7 studies to support their contention that circumcision may lead to sexual problems in circumcised men and their partners.

Four of the studies involved only men circumcised as adults<sup>11–14</sup> or some men circumcised as adults.<sup>15</sup> Men circumcised as adults most frequently have the procedure performed for medical reasons, which introduces both physical and psychological factors that may affect their reporting of sexual difficulties. In 2 of the studies cited by Frisch et al, a significant number of men reported improved satisfaction after circumcision.<sup>11,13</sup> Interestingly, another of their cited studies<sup>14</sup> concluded that circumcision had neither a negative nor a positive effect on the female partner’s perception of sexual satisfaction, a conclusion that contradicts that of Frisch et al. **The shortcomings of the study by Sorreels et al<sup>16</sup> are discussed in our technical report.** Finally, the study by Frisch et al used a cross-sectional survey of Danish men that found that circumcised men were more likely to report sexual difficulties than **uncircumcised** men.<sup>17</sup> Circumcised men represented only 5% of 2343 sexually experienced survey respondents, and only 15% of those circumcised men ( $n = 17$ ) had the procedure in the first 6 months of life. Attributing these findings to decreased penile sensitivity is a stretch. It seems far more likely that the findings are attributable to the kinds of social bias the authors attribute to the AAP task force. Male circumcision is rare in Denmark, rare enough that circumcised males are epidemiologic outliers, which may lead some of them to feel “different,” leading to anxiety about sexual experiences with women who perceive a circumcised penis as abnormal. In addition, because many of the circumcised men had the procedure performed later in life, some likely for medical reasons, they are far from representative of a group of men circumcised as infants. It should be noted that the findings of the Danish survey contrast starkly to

those of 2 randomized controlled trials from Africa, which are discussed in the AAP technical report.<sup>18,19</sup>

## AGE AT CIRCUMCISION

A central claim of Frisch et al is that if circumcision is to be performed before an age at which an individual can decide for himself, there must be a compelling reason for doing so. They argue that there is no compelling reason for performing a circumcision before sexual debut and additionally claim that “sexually transmitted HIV infection is not a relevant threat to children.”

Underlying the authors' views are several presuppositions that reflect the ideal, but not the reality, of human decision making. The first of these is that the responsible use of condoms will “provide close to 100% reduction in risk for any STIs.” We agree, and fully support efforts to make sexual activity as safe as possible through the routine use of condoms. However, despite huge educational efforts, many individuals around the world do not use condoms consistently. If they did, sexually transmitted infection and HIV would decline to the point of nonexistence. The added protective benefit of circumcision exists precisely because responsible condom use is far from universal. In 2011, 39.8% of sexually active high school students in the United States reported not using a condom during their last sexual intercourse.<sup>20</sup> Condoms represent one of several tools for reducing the risk of infections transmitted during sexual contact. Circumcision is another.

The claim that “sexually transmitted HIV infection is not a relevant threat to children” is incorrect given the US data. In 2011, sexual debut occurred at or before age 13 years for 6.2% of US high school students, and **most people in the United States are sexually active before the age at which they would possess**

**the legal authority to consent to a circumcision.**<sup>20</sup> A nationwide sample of adolescent females between the ages of 14 and 19 years estimated that 4 in 10 sexually active adolescent females have a sexually transmitted infection (defined as human papillomavirus, chlamydia, trichomoniasis, genital herpes virus, or gonorrhea).<sup>21</sup> National HIV surveillance data show that, in 2008, there were at least 2266 HIV infections among US adolescents 13 to 19 years of age.<sup>22</sup>

## ETHICAL ISSUES

Frisch et al claim that “the AAP report lacks a serious discussion of the central ethical dilemma with, on one side, parents' right to act in the best interest of the child ... and, on the other side, infant boys' basic right to physical integrity in the absence of compelling reasons for surgery.” **The authors' argument about the basic right to physical integrity is an important one,** but it needs to be balanced by other considerations. The right to physical integrity is easier to defend in the context of a procedure that offers no potential benefit, but the assertion by Frisch et al of ‘no benefit’ is clearly contradicted by the published scientific peer-reviewed evidence. Although task force members did not find the data sufficiently compelling to justify a recommendation for routine neonatal circumcision, **we did find that the benefits are substantial enough to allow parents to make this decision for their male children.** This stance is **very similar** to that of The **Canadian Medical Society,** the British Medical Association, and the Royal Australasian College of Physicians. Frisch et al appeal to the ethical precept “First, do no harm,” but they fail to recognize that in situations in which a preventive benefit exists, harm can also be done by failing to act. Whereas there are rare

situations in which a male will be harmed by a circumcision procedure, it is also true that **some males will be harmed by not being circumcised.** Simply because it is difficult to identify exactly which individuals have suffered a harm because they were not circumcised should not lead one to discount the very real harms that might befall some men by not being circumcised. There is no easy answer to this issue ethically. Regardless of what decision is made on behalf of a young male, harm might result from that decision. That is precisely why the AAP task force members found that this decision properly remains with parents and that parents should have information about both potential benefits and potential harms as they make this decision for their child.

## TASK FORCE ON CIRCUMCISION 2012

Susan Blank, MD, MPH, Chairperson  
Michael Brady, MD, Representing the AAP Committee on Pediatrics AIDS and Committee on Infectious Disease  
Ellen Buerk, MD, Representing the AAP Board of Directors

Waldemar Carlo, MD, Representing the AAP Committee on Fetus and Newborn  
Douglas Diekema, MD, MPH, Representing the AAP Committee on Bioethics

**Andrew Freedman,** MD, Representing the AAP Section on Urology  
Lynne Maxwell, MD, Representing the AAP Section on Anesthesiology  
Steven Wegner, MD, JD, Representing the AAP Committee on Child Health Financing

## LIAISONS

Charles LeBaron, MD – *Centers for Disease Control and Prevention*

Lesley Atwood, MD – *American Academy of Family Physicians*

Sabrina Craigo, MD – *American College of Obstetricians and Gynecologists*

## REFERENCES

1. American Academy of Pediatrics Task Force on Circumcision. Male circumcision. *Pediatrics*. 2012;130(3). Available at: [www.pediatrics.org/cgi/content/full/130/3/e756](http://www.pediatrics.org/cgi/content/full/130/3/e756)
2. Cummins RO, Hazinski MF. The most important changes in the international ECC and CPR guidelines 2000 [editorial]. *Circulation*. 2000;102(8 suppl):I371–I376
3. World Health Organization; UNAIDS. New data on male circumcision and HIV prevention: policy and programme implications. Geneva, Switzerland: World Health Organization; 2007. Available at: [http://libdoc.who.int/publications/2007/9789241595988\\_eng.pdf](http://libdoc.who.int/publications/2007/9789241595988_eng.pdf). Accessed January 7, 2013
4. Moldwin RM, Valderrama E. Immunohistochemical analysis of nerve distribution patterns within prepuccial tissue (abstract). *J Urol*. 1989;141(4 pt 2):499A
5. Taylor JR, Lockwood AP, Taylor AJ. The prepuce: specialized mucosa of the penis and its loss to circumcision. *Br J Urol*. 1996;77(2):291–295
6. Tuncali D, Bingül F, Talim B, Surucu S, Sahin F, Aslan G. Histologic characteristics of the human prepuce pertaining to its clinical behavior as a dual graft. *Ann Plast Surg*. 2005;54(2):191–195
7. Wu ZM, Chen YF, Qiu PN, Ling SC. Correlation between the distribution of SP and CGRP immunopositive neurons in dorsal root ganglia and the afferent sensation of preputial frenulum. *Anat Rec (Hoboken)*. 2011;294(3):479–486
8. Podnar S. Clinical elicitation of the peniloscavernosus reflex in circumcised men. *BJU Int*. 2012;109(4):582–585
9. Smith DK, Taylor A, Kilmarx PH, et al. Male circumcision in the United States for the prevention of HIV infection and other adverse health outcomes: report from a CDC consultation. *Public Health Rep*. 2010;125 (suppl 1):72–82
10. Yang DM, Lin H, Zhang B, Guo W. Circumcision affects glans penis vibration perception threshold [in Chinese]. *Zhonghua Nan Ke Xue*. 2008;14(4):328–330
11. Fink KS, Carson CC, DeVellis RF. Adult circumcision outcomes study: effect on erectile function, penile sensitivity, sexual activity and satisfaction. *J Urol*. 2002;167 (5):2113–2116
12. Kim D, Pang MG. The effect of male circumcision on sexuality. *BJU Int*. 2007;99(3): 619–622
13. Shen Z, Chen S, Zhu C, Wan Q, Chen Z. Erectile function evaluation after adult circumcision [in Chinese]. *Zhonghua Nan Ke Xue*. 2004;10(1):18–19
14. Cortés-González JR, Arratia-Maqueo JA, Gómez-Guerra LS. Does circumcision have an effect on female's perception of sexual satisfaction? [in Spanish]. *Rev Invest Clin*. 2008;60(3):227–230
15. Tang WS, Khoo EM. Prevalence and correlates of premature ejaculation in a primary care setting: a preliminary cross-sectional study. *J Sex Med*. 2011;8(7):2071–2078
16. Sorrells ML, Snyder JL, Reiss MD, et al. Fine-touch pressure thresholds in the adult penis. *BJU Int*. 2007;99(4):864–869
17. Frisch M, Lindholm M, Grønbaek M. Male circumcision and sexual function in men and women: a survey-based, cross-sectional study in Denmark. *Int J Epidemiol*. 2011;40(5):1367–1381
18. Kigozi G, Watya S, Polis CB, et al. The effect of male circumcision on sexual satisfaction and function, results from a randomized trial of male circumcision for human immunodeficiency virus prevention, Rakai, Uganda. *BJU Int*. 2008;101(1):65–70
19. Krieger JN, Mehta SD, Bailey RC, et al. Adult male circumcision: effects on sexual function and sexual satisfaction in Kisumu, Kenya. *J Sex Med*. 2008;5(11):2610–2622
20. Centers for Disease Control and Prevention. 1991-2011 High school youth risk behavior survey data. Available at: <http://apps.nccd.cdc.gov/youthonline>. Accessed January 7, 2013
21. Forhan SE, Gottlieb SL, Sternberg MR, et al. Prevalence of sexually transmitted infections among female adolescents aged 14 to 19 in the United States. *Pediatrics*. 2009;124 (6):1505–1512
22. Centers for Disease Control and Prevention. HIV surveillance in adolescents and young adults. Available at: [www.cdc.gov/hiv/topics/surveillance/resources/slides/adolescents/slides/Adolescents.pdf](http://www.cdc.gov/hiv/topics/surveillance/resources/slides/adolescents/slides/Adolescents.pdf). Accessed January 7, 2013

**Cultural Bias and Circumcision: The AAP Task Force on Circumcision  
Responds**

TASK FORCE ON CIRCUMCISION

*Pediatrics*; originally published online March 18, 2013;

DOI: 10.1542/peds.2013-0081

<b>Updated Information &amp; Services</b>	including high resolution figures, can be found at: <a href="http://pediatrics.aappublications.org/content/early/2013/03/12/peds.2013-0081.citation">http://pediatrics.aappublications.org/content/early/2013/03/12/peds.2013-0081.citation</a>
<b>Citations</b>	This article has been cited by 1 HighWire-hosted articles: <a href="http://pediatrics.aappublications.org/content/early/2013/03/12/peds.2013-0081.citation#related-urls">http://pediatrics.aappublications.org/content/early/2013/03/12/peds.2013-0081.citation#related-urls</a>
<b>Permissions &amp; Licensing</b>	Information about reproducing this article in parts (figures, tables) or in its entirety can be found online at: <a href="http://pediatrics.aappublications.org/site/misc/Permissions.xhtml">http://pediatrics.aappublications.org/site/misc/Permissions.xhtml</a>
<b>Reprints</b>	Information about ordering reprints can be found online: <a href="http://pediatrics.aappublications.org/site/misc/reprints.xhtml">http://pediatrics.aappublications.org/site/misc/reprints.xhtml</a>

PEDIATRICS is the official journal of the American Academy of Pediatrics. A monthly publication, it has been published continuously since 1948. PEDIATRICS is owned, published, and trademarked by the American Academy of Pediatrics, 141 Northwest Point Boulevard, Elk Grove Village, Illinois, 60007. Copyright © 2013 by the American Academy of Pediatrics. All rights reserved. Print ISSN: 0031-4005. Online ISSN: 1098-4275.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™

